

Intravitreal Axitinib Hydrogel (OTX-TKI) in Neovascular Age-Related Macular Degeneration: Efficacy and Safety Results from the Phase 3 SOL-1 Trial

Dilsher S. Dhoot, MD

California Retina Consultants, Santa Barbara, CA

Arshad M. Khanani, Patricio G. Schlottmann, Mark R. Barakat, Jeffrey S. Heier, Peter K. Kaiser, and Nadia K. Waheed on behalf of SOL-1 study group

The following presentation discusses OTX-TKI - an investigational product candidate that has not been approved by the FDA or any other regulatory body; safety and effectiveness have not been established.

Financial Disclosures (Dilsher S. Dhoot): Consultant: 4DMT, Abbvie, Adverum, Alcon Pharmaceuticals, Alimera Sciences, Inc., Alkeus, Allergan, ANI Pharmaceuticals, Amgen, Annexon, Apellis Pharmaceuticals, Inc., Astellas; Bausch and Lomb, Bayer Healthcare Pharmaceuticals, Inc., Biocryst, Coherus, EyePoint Pharmaceuticals, Genentech, Harrow, IvericBio, Novartis, Ocular Therapeutix, Oculis, Optos, Inc., Opthea, Outlook Therapeutics, Oxular, Regeneron, REGENXBIO, RetinaAI, Roche, Sanofi, Santen, Inc. **Research Support:** 4DMT, Adverum, Annexon Biosciences, Eyebio, Eyepoint Pharma, Genentech, Kodiak Sciences, Ocular Therapeutix, Inc, Oculis, Opthea, Regeneron, RegenxBio, Sanofi. **Speaker:** Amgen, ANI Pharmaceuticals, Apellis, Genentech, Regeneron

Study Design: OTX-TKI and Aflibercept Direct Comparison


Key Enrollment Criteria

- Treatment naïve
- **BCVA ≥ 54 ETDRS letters** (~20/80)
- **CSFT of ≤ 500 μm**

Key Randomization Criteria

- **BCVA gain ≥ 10 ETDRS letters**
OR
- **BCVA ≥ 84 ETDRS letters** (~20/20)
- **CSFT of ≤ 350 μm**

Proportion of subjects who maintained visual acuity, defined as <15 ETDRS letters of BCVA loss from baseline at Week 36



Baseline (Randomization) Study Eye Characteristics

Mean (SD) BCVA, ETDRS letter score,
Snellen Equivalent

OTX-TKI 0.45 mg
N = 172[†]

80.8 (7.6)
~20/25

Aflibercept 2 mg
N = 172

79.2 (7.9)
~20/25

Mean (SD) CSFT, μm ,

219.3 (37.1)

226.8 (42.1)

*No aflibercept injection is given at baseline; [†]Two OTX-TKI subjects were misrandomized, did not receive study treatment and were not included in the full analysis set

ClinicalTrials.gov. identifier: NCT06223958

Rescue Treatment Criteria: BCVA loss ≥ 15 ETDRS letters from baseline due to nAMD; or New macular hemorrhage that would likely lead to irreversible vision loss if left untreated in the opinion of the investigator after discussion with Medical Monitor


BCVA, best-corrected visual acuity; CSFT, central subfield thickness; ETDRS, Early Treatment Diabetic Retinopathy Study; SD, standard deviation

Unmatched Durability: First Trial to Demonstrate Durability >9 Months

Maintenance of Visual Acuity*


Primary Endpoint

Key Secondary Endpoint



Time to First Rescue Treatment

Key Secondary Endpoint



■ OTX-TKI 0.45 mg (n=170) ■ Aflibercept 2 mg (n=172)

*Maintenance of vision defined as proportion of subjects who maintain visual acuity, a loss of <15 ETDRS letters of BCVA from baseline

†Nominal P value

Predefined statistical rules were applied to adjust for treatment discontinuation or deviation as per the pre-specified statistical analysis plan. Full analysis set. Rescue-free subjects include those who did not require rescue injections as specified by the protocol.


BCVA, best-corrected visual acuity; ETDRS, Early Treatment Diabetic Retinopathy Study

Sustained Disease Control: Visual and Anatomic Stability with OTX-TKI

Mean Change in BCVA and CSFT from Screening


Post Hoc Analysis

Primary Outcome



Mean Change in Retinal Fluid Volume

Prespecified Exploratory Endpoint



Full analysis set. For subjects who received any rescue injections, their observed BCVA and CSFT values after the 1st rescue injection are set to missing. Baseline measurements are the last non-missing measurement prior to the study drug injection on Day 1. Error bars represent standard error
AFL, aflibercept; BCVA, best-corrected visual acuity; CSFT, central subfield thickness, measured from the internal limiting membrane to the retinal pigment epithelium; ETDRS, Early Treatment Diabetic Retinopathy Study; IRF, intraretinal fluid; SRF, subretinal fluid

OTX-TKI was Generally Well Tolerated

Subjects with Ocular AEs in the Study Eye (>2%) Through Week 52, n (%)	OTX-TKI 0.45 mg n = 170	Aflibercept 2 mg n = 172
Vitreous floaters	21 (12.4)	2 (1.2)
Cataract	12 (7.1)	5 (2.9)
Conjunctival hemorrhage	11 (6.5)	5 (2.9)
Retinal hemorrhage	10 (5.9)	17 (9.9)
Dry eye	7 (4.1)	2 (1.2)
Vitreous detachment	7 (4.1)	3 (1.7)
Punctate keratitis	6 (3.5)	0
Vitreous opacities	6 (3.5)	0
Eye pain	5 (2.9)	1 (0.6)
Anterior chamber opacity	4 (2.4)	0
Posterior capsule opacification	4 (2.4)	6 (3.5)

Vitreous Floaters

- All cases were mild to moderate in severity
- No evidence of associated IOI
- Onset coincides with drug elution stage of OTX-TKI hydrogel bioresorption process
- For all subjects with vitreous floaters AE reported, drug particles are no longer visible

Intraocular Inflammation

- Nine events observed in study eyes of 7 subjects in the OTX-TKI arm
 - All cases were mild or moderate in severity and resolved
- **There were no cases of endophthalmitis, occlusive or non-occlusive retinal vasculitis**